

PROBLEMS IN DIATOM ANALYSIS OF DEPOSITS: ALLOCHTONOUS VALVES AND FRAGMENTATION¹

L. BEYENS² & L. DENYS³

ABSTRACT

Beyens, L. & L. Denys 1982 Problems in diatom analysis of deposits: Allochtonous valves and fragmentation – Geol. Mijnbouw 61: 159-162.

A review of the various sources of allochtonous diatom valves in deposits is given. The importance of these problems in palaeoecological research is discussed and some simple methods for their approach in quantitative analysis are introduced.

INTRODUCTION

Selective dissolution, fragmentation and influx of allochtonous valves are serious problems encountered in quantitative diatom-analysis of sediment cores for palaeoecological purposes, the effects of these phenomena being such that they have to be taken into account for each interpretation of obtained data. The process and consequences of selective dissolution have already been treated extensively (BRADBURY, 1973; JOHNSON, 1974; JØRGENSEN, 1955; KAMATANI, 1969; KAMATANI & RILEY, 1979; LEWIN, 1961; ROUND, 1964; SCHRADER, 1971). Influx of allochtonous valves and fragmentation, however, have been somewhat neglected until now.

ALLOCHTONOUS VALVES

Due to their small dimensions and their particular shape, living diatoms and their valves are easily transported by moving water and air. For this reason, valves not belonging to the autochtonous flora are frequently found in deposits. It is obvious that it is of the greatest importance to differentiate between these allochtonous valves and those belonging to the original assemblage, for only the latter can give precise information on the local palaeoenvironment.

Sources

Important transport of diatoms and their remnants can occur lateral as well as vertical. Lateral transport by marine currents may involve enormous distances (BURCKLE, 1978). Transport in watercourses and run-off (HENDEY, 1974) result in an important seaward flux of fresh-water taxa, while marine and brackish taxa are carried far up into estuaries by tidal currents (BROCKMANN, 1940; GESSNER & SIMONSEN, 1967; SIMONSEN, 1969). In some cases, the allochtonous population can even exceed the autochtonous one quantitatively as well as qualitatively (SIMONSEN, 1969), which may easily lead to erroneous conclusions.

Allochtonous valves in lake deposits can be brought in from outside of the sedimentary basin, but also from different places in the lake itself. Solifluction (EVANS, 1970) and seasonal water movements (MERILAINEN, 1969) can lead to the deposition of littoral forms in the more profound parts of a lake, thus destroying the expected horizontal distribution of the valves in the bottom sediments. Aerophilous forms may be used to indicate the washing in of material (WALKER, 1978). Some caution should, however, be made in applying this assumption, since most aerophilous taxa may thrive in purely aquatic conditions as well.

Lateral transport by moving air (VAN OVEREEM, 1937; GEISSLER & GERLOFF, 1966; KALBE, 1973) can also be of some importance. DU SAAR (1967) investigated the transportation of living diatoms and valves by a strong seabreeze, and concluded that living diatoms (mainly fragile planktonic species) were blown on land in spray-droplets, while valves of strongly silicified benthonic forms were taken up from the beach sand and transported over many metres. Animals, for instance

¹ Manuscript received: 1981-09-04. Revised manuscript accepted: 1981-11-28.

² Dienst Algemene Plantkunde, RUCA Groenenborgerlaan 171, B-2020 Antwerpen, Belgium.

³ Zilverboslaan 8. B-2070 Ekeren, Belgium.

foraging birds (SIMONSEN, 1957), and human activities (ships, etc.) are also possibilities for lateral transport.

Vertical transport of valves in deposits is still badly known. Possible causes are reworking and infiltration. Both upward and downward movements occur. Probably, infiltration-effects can be quite important in coarse sediments involving small valves. Percolating rain- and melting water as well as fluctuations of the ground water level may displace valves. Fresh-water deposits may become contaminated with marine diatoms during a flooding with seawater as a result of infiltration and wave action. Vertical displacement may also result from bioturbation.

Detection

It is often difficult to recognise the allochthonous component of a fossil assemblage, although it may form an important part of it. SIMONSEN (1969) suggested that benthonic forms should be used in order to find out the autochthonous component, since planctonic taxa are subjected to transport more frequently. Rare taxa, present in large numbers may also be regarded as autochthonous (BROCKMANN, 1940). A further discrimination may be based on ecological data. Sometimes the presence of rheophilous forms can be linked with lateral transport phenomena. A high degree of fragmentation does not necessarily coincide with transport over a long distance and should thus not be considered a very useful criterium. Recognition of vertical transport is often only possible when other (lithostratigraphic or paleontological) evidence is present as in the example given in Fig. 1.




Fig. 1
Part of a section, taken in a bog at I-penrooi (Belgian Campine) showing a hiatus between 55 and 53 cm. (Beyens, 1980). An abrupt change in the vegetation (Arboreal/Non-Arboreal pollen) is observed at the same level, while the corresponding change in the diatom flora is situated beneath this boundary. This indicates that there has been a vertical migration of the *T. flocculosa* valves, which affected the composition of the assemblage at 55 cm in such a way that the percentage of *Frustulia*, which is in fact the most important autochthonous species at this level, is reduced to an insignificant value.

Quantifying the allochthonous component

Knowledge of the amount in which allochthonous valves are present in analysed samples can give valuable indications on the physical and hydrographical environment during deposition and can simplify the interpretation of the data obtained. We can try to quantify the allochthonous component in such a way that subsequent samples can be compared mutually.

The following method uses salinity dependence and benthonic life style to differentiate allochthonous valves from autochthonous ones. It should, however, be mentioned that other ecological requirements (pH, trophic conditions, . . .) can be used in the same way. The salinity group in the classification of VAN DER WERFF & HULS (1957-1974) with the largest percentage of benthonic forms, the 'optimal group', is calculated for each sample. The percentage of this group is added to those of the two 'neighbour groups', giving a good estimation of the autochthonous group. Next, the percentages of the allochthonous groups on the fresh-water side and on the marine side of the autochthonous group are calculated and expressed as ratios (F- and M-values) of this group. An exemplary calculation follows:

optimal group	brackish	30%	
neighbour groups:	brackish-fresh	15%	
	brackish-marine	10%	
	Σ	55%	autochthonous component
fresh-water allochthonous group:	fresh-brackish	5%	
	fresh	1%	
	Σ	6%	F-value: 0,11
marine allochthonous group:	marine-brackish	20%	
	marine	19%	
	Σ	39%	M-value: 0,71

Graphic figuration of these values, as in Fig. 2, gives a clear picture of the changes in the influx of allochthonous valves.

FRAGMENTATION

One of the most annoying disturbances, when counting a diatom slide, is the presence of broken valves. It is only in some more recent papers (e.g. CLARYSSE, 1974; PALS, ET AL., 1980; VOORRIPS & JANSMA, 1974) that this evil has been faced, and taken into account when interpreting the data.

Causes

Fragmentation by abrasion can result from various types of transport. This transport does not necessarily occur over considerable geographic distances. As a result of repeated resuspension by wave- or tidal action, a valve may be transported over a very long distance and still not be embedded far from its

- calculate the 'mean number' of complete valves (for valves with a more strongly developed apical axis) by means of the formula:

$$N = \frac{n\alpha + n}{2}$$

with n: the number of fragments

$\alpha = 1/L$, being the mean length of the fragments,

L being the mean length of the complete valves.

Since α is species and sample specific, the last method can be very time-consuming. In addition, following difficulties may arise:

- if not enough complete valves are present to calculate a meaningful value of L, a similar value must be derived from literature data or if possible from a known length/width ratio
- larger valves break more rapidly than small ones so L will be underestimated
- fragments cannot always be identified.

Fig. 3 illustrates the difference in the quantitative importance of *Pinnularia viridis* in a holocene bog section, between inclusion and exclusion of fragments in the sum by means of the 'mean number' method.

Although the inclusion of fragments in the sum is time consuming and always loaded with errors, its advantages may be worthwhile for any palaeoecological interpretation of diatom data.

REFERENCES

- Berglund, B. E. 1971 Littorina transgressions in Blekinge, South Sweden, a preliminary survey – Geologiska Föreningen Stockholm Förhandlingar 93: 626-652.
- Beyens, L. 1980 Diatom, pollen and rhizopod analyses on a Subatlantic peat bog section from Ipenrooi (North Belgium) – Abstr. 5th. Int. Palyn. CXONF.: 40.
- Bradbury, J. P. 1973 Ecology of freshwater diatoms – Nova Hedwigia XXIV: 145-168.
- Brockmann, C. 1940 Diatomeen als Leitfossilien in Küstenablagerungen – Westküste 2: 150-181.
- Burckle, L. H. 1978 Marine Diatoms. In B. U. Haq & A. Boersma (ed.) – Introduction to marine micropaleontology. Elsevier, North-Holland, Inc. (New York): 245-266.
- Clarysse, R. 1974 The diatom flora of the Steenbrugge-clay (Eemian). In: J. Vandenberghe, N. Vandenberghe & F. Gullentops: Late Pleistocene and Holocene in the neighbourhood of Brugge – Med. Kon. Acad. Wet. België. Klasse Wet. XXXVI: 72-77.
- Denys, L. 1981 Diatomeeënstudie van de Holocene afzettingen in de Belgische kustvlakte – Licenciaatsverh. UIA: 130 pp.
- Du Saar, A. 1967 Results and problems in diatom investigation at the Geological Survey in Haarlem – Geol. Mijnbouw 46: 105-108.
- Evans, G. H. 1970 Pollen and diatom analysis of Late-Quaternary deposits in the Blelham basin, North Lancashire – New Phytologist 69: 821-874.
- Geissler U. & J. Gerloff 1966 Das Vorkommen von Diatomeen in menschlichen Organen und in der Luft – Nova Hedwigia 10: 565-577.
- Gessner, F. & R. Simonsen 1967 Marine diatoms in the Amazon? – Limnol. Oceanogr. 12: 709-711.
- Hendey, N. I. 1974 A revised check-list of British marine diatoms – J. Mar. Biol. Ass. U.K. 54: 277-300.
- Johnson, T. L. 1974 The dissolution of siliceous microfossils in surface sediments of the eastern tropical Pacific – Deep-Sea Res. 21: 851-864.
- Jørgensen, E. G. 1955 Solubility of the silica in diatoms – Physiol. Plantarum 8: 846-851.
- Kalbe, L. 1973 Kieselalgen in Binnengewässern – Ziemsen Verlag (Wittenberg): 206 pp.
- Kamatani, A. 1969 Regeneration of organic nutrients from diatom decomposition – J. Oceanogr. Soc. Japan 25: 63-74.
- Kamatani, A. & J. P. Riley 1979 Rate of dissolution of diatom silica walls in seawater – Mar. Biology 55 1: 29-35.
- Lewin, J. C. 1961 The dissolution of silica from diatom walls – Geochim. Cosmochim. Acta 21: 182-198.
- Meriläinen, J. 1969 Distribution of diatom frustules in recent sediments of some meromictic lakes – Mitt. Internat. Verein. Limnol. 17: 186-192.
- Pals, J. P., B. Van Geel & A. Delfos 1980 Palaeoecological studies in the Klokkeweel Bog near Hoogkarspel (prov. of Noord Holland) – Rev. Palaeobot. Palynol. 30: 371-418.
- Round, F. E. 1964 The diatom sequence in lake deposits: some problems of interpretation – Verh. Internat. Verein. Limnol. 15: 1012-1020.
- Schrader, H. J. 1971X Fecal pellets: role in sedimentation of pelagic diatoms – Science 174: 55-57.
- Simonsen, R. 1957 Spätglaziale Diatomeen aus Holstein – Arch. f. Hydrobiol. 53: 337-349.
- Simonsen, R. 1969 Diatoms as indicators in estuarine environments – Veröffentl. Inst. Meeresforsch. Bremerhaven 11: 287-291.
- Van der Werff, A. & H. Huls 1957-1974 Diatomeeënflora van Nederland. Reprint 1976 – Otto Koeltz Science Publ. (Koenigstein):
- Van Overeem, M. A. 1937 On green organisms occurring in the lower troposphere – Rec. Trav. Bot. Neerl. 34: 388-442.
- Voorrips, A. & M. J. Jansma 1974 Pollen and diatom analysis of a shore section of the former lake Wervershoof – Geol. Mijnbouw 53: 429-435.
- Walker, R. 1978 Diatom and pollen studies of a sediment profile from Melynllyn, a mountain tarn in Snowdonia, North Wales – New Phytologist 81: 791-804.